

Søgeprotokol for Nationale Kliniske Retningslinjer

Projekttitel/aspekt	NKR for medicinsk behandling af bipolar lidelse - Baggrundssøgning
Kontakt projektgruppe	Stine Ulendorf Jacobsen
Kontakt bibliotek	Hanne Munch Kristiansen, AU Library, Psykiatri
Refworks login	
Vedhæftede filer	
Senest opdateret	11. august 2013 (BHP)

Baggrund	<p>Sundhedsstyrelsen igangsætter i august 2013 arbejdet med nationale kliniske retningslinjer for medicinsk behandling af bipolar lidelse.</p> <p>Ca. 1 % af den danske befolkning svarende til omkring 40.000 mennesker lider af bipolar lidelse. Bipolar lidelse er karakteriseret ved hyppige depressive og maniske episoder med forværring i løbet af sygdommen, hyppige psykiatriske indlæggelser, stor risiko for nedsat arbejdsevne og førtidspensionering samt 20 gange øget selvmordsrisiko i forhold til baggrundsbefolkningen.</p> <p>Behandlingen af bipolar lidelse består af kombineret psykofarmakologisk og psykologisk behandling (psykoedukation). Den farmakologiske behandling er yderst kompleks, idet den dels afhænger af, hvilken fase patienten befinder sig i (mani, depression, remission), dels fordi kombinationsbehandling fra forskellige farmakologiske klasser er nødvendig ved 90 % af patienterne (lithium, anti-epileptika, antipsykotika og muligvis antidepressiva).</p> <p>Der er stor variation i den farmakologiske behandling af bipolar lidelse dels mellem forskellige regioner i Danmark, såvel under ind-læggelse som i ambulant regi, og dels blandt privatpraktiserende psykiatere og praktiserende læger.</p>
søgetermer	<p>Engelsk: Bipolar disorder; bipolar affective disorder; manic depression; manic-depressive disorder</p> <p>Dansk : Bipolar lidelse, mani-depressiv sygdom</p> <p>Norsk: bipolar lidelse, bipolar sykdom ??</p> <p>Svensk: bipolar lidelse, bipolar sjukdom??</p>
Inklusions- og eksklusionskriterier	<p>Sprog: Engelsk, dansk, norsk og svensk</p> <p>År: de sidste 10 år, (2002 -2013)</p> <p>Population: Alle</p> <p>Publikationstyper: Guidelines, practice guidelines, clinical guidelines, MTV, systematiske reviews</p>

Informationskilder

Database	Interface	Dato for søgning
G-I-N International	Internet	17.06.2013
NICE (UK)	Internet	17.06.2013
National Guideline Clearinghouse (USA)	Internet	17.06. 2013
Scottish Intercollegiate Guidelines Network (SIGN)	Internet	18.06.2013
HTA Databasen	Internet	21.06.2013
The Cochrane Library	Internet	21.06.2013
SBU, Sverige	Internet	18.06.2013
Socialstyrelsen, Sverige	Internet	18.06.2013
Helsedirektoratet, Norge	Internet	18.06.2013
Kunnskapscentret, Norge	Internet	18.06.2013

Database	Interface	Dato for søgning
PubMed	Internet	18.06.2013
EMBASE	Embase.com Elsevier	19.06.2013
CINAHL	Ebsco	20.06.2013
PSYCINFO	ProQuest	20.06.2013

Note: Subject headings og inklusions- og eksklusionskriterier er tilpasset de enkelte databaser.

Søgestrategi

G-I-N

21 fund, 8 relevante

1.	Bipolar affective disorder (SIGN CPG 82)	SIGN (GB) - Scottish Intercollegiate Guidelines Network	Guideline	Jul 01, 2005	United Kingdom	Published
2.	Bipolar disorder (CG38)	NICE (UK) - National Institute for Health and Care Excellence	Guideline	Jul 01, 2012	United Kingdom	Under review
3.	Bipolar Disorder	MoH (SG) - Ministry of Health, Singapore	Guideline	Dec 01, 2011	Singapore	Published
4.	Kaksisuuntainen mielialahäiriö [Bipolar affective disorder]	CC (FI) - Current Care / Duodecim - Finnish Medical Society	Guideline	Oct 28, 2008	Finland	Published
5.	Troubles bipolaires. Guide ALD [Bipolar disorder]	HAS (FR) - French National Authority for Health (formerly: ANAES)	Guideline	May 01, 2009	France	Published
6.	VA/DoD clinical practice guideline for management of bipolar disorder in adults. Department of Defense. Department of Veterans Affairs. Veterans Health Administration. NGC:007865	AHRQ (US) - Agency for Healthcare Research and Quality	Guideline Clearing Report	May 01, 2010	United States	Published
7.	Bipolare Störungen. S3-LL (DGPPN, DGBS) [Bipolar disorders]	AWMF (DE) - Association of Scientific Medical Societies	Guideline	May 11, 2012	Germany	Published
8.	Retningslinjer for behandling av bipolare lidelser	Hdir (NO) - Norwegian	Guideline	Nov 01,	Norway	In development


NICE

“bipolar disorder” , clinical guidelines 21 fund, 2 relevante

1. **Bipolar disorder (CG38)**

The summary of the published clinical guideline on **Bipolar disorder** . It links to the published guidance and key documents.

[View the summary and implementation tools](#) 26 July 2006

-
- [Full Guidance](#)
 - [Quick reference guide](#)
 - [NICE Guidance](#)
 -  [Information for the public](#)
 -

2. **Referred Bipolar disorder (update)**

The summary of the referred clinical guideline on **Bipolar disorder (update)**. It links to the referred guidance and key documents.

National Guideline Clearinghouse

Guidelines by Topic

[Disease/Condition](#) > [Psychiatry and Psychology](#) (456) > [Mental Disorders](#) (313) > [Mood Disorders](#) (62) > [Affective Disorders, Psychotic](#) (12) >

Bipolar Disorder (11)

1	11
---	----

 1-11 of 11

- | | | |
|----|---|--|
| 1. | Occupational therapy practice guidelines for adults with serious mental illness. 2012. NGC:009369
American Occupational Therapy Association, Inc. - Professional Association. View all guidelines by the developer(s) | Compare
Selected <input type="checkbox"/>
38528
0 |
| 2. | Bipolar disorder. 2011 Nov. NGC:009526
Singapore Ministry of Health - National Government Agency [Non-U.S.]. View all guidelines by the developer(s) | Compare
Selected <input type="checkbox"/>
39325
0 |
| 3. | Depression. 1998 Jun (revised 2011 Aug). NGC:008672
University of Michigan Health System - Academic Institution. View all guidelines by the developer(s) | Compare
Selected <input type="checkbox"/>
34406
0 |
| 4. | Depression and mania in patients with HIV/AIDS. 2005 Feb (revised 2010 Oct). NGC:008645
New York State Department of Health - State/Local Government Agency [U.S.]. View all guidelines by the developer(s) | Compare
Selected <input type="checkbox"/>
34265
0 |
| 5. | VA/DoD clinical practice guideline for management of bipolar disorder in adults. 2010 May. NGC:007865
Department of Defense - Federal Government Agency [U.S.]; Department of Veterans Affairs - Federal Government Agency [U.S.]; Veterans Health Administration - Federal Government Agency [U.S.]. View all guidelines by the developer(s) | Compare
Selected <input type="checkbox"/>
16314
0 |

- | | | | |
|--|---|-------|---|
| <p>6. Anxiety and depression in children and youth—diagnosis and treatment. 2010
 Jan 1. NGC:009471
 Medical Services Commission, British Columbia - State/Local Government Agency
 [Non-U.S.]. View all guidelines by the developer(s)</p> | <p>Compare</p> <p>Selected <input type="checkbox"/></p> <table border="1" style="width: 100%; border-collapse: collapse;"> <tr><td style="text-align: center;">38904</td></tr> <tr><td style="text-align: center;">0</td></tr> </table> | 38904 | 0 |
| 38904 | | | |
| 0 | | | |
| <p>7. Clinical practice guidelines for psychosocial interventions in severe mental illness. 2009. NGC:008509
 Aragon Institute of Health Sciences-I+CS - State/Local Government Agency [Non-U.S.]; GuiaSalud - National Government Agency [Non-U.S.]; Ministry of Health (Spain) - National Government Agency [Non-U.S.]. View all guidelines by the developer(s)</p> | <p>Compare</p> <p>Selected <input type="checkbox"/></p> <table border="1" style="width: 100%; border-collapse: collapse;"> <tr><td style="text-align: center;">33135</td></tr> <tr><td style="text-align: center;">0</td></tr> </table> | 33135 | 0 |
| 33135 | | | |
| 0 | | | |
| <p>8. Use of psychiatric medications during pregnancy and lactation. 2008 Apr.
 NGC:006421
 American College of Obstetricians and Gynecologists - Medical Specialty Society. View all guidelines by the developer(s)</p> | <p>Compare</p> <p>Selected <input type="checkbox"/></p> <table border="1" style="width: 100%; border-collapse: collapse;"> <tr><td style="text-align: center;">12490</td></tr> <tr><td style="text-align: center;">0</td></tr> </table> | 12490 | 0 |
| 12490 | | | |
| 0 | | | |
| <p>9. Mental health disorders among substance-using HIV-infected patients. 2008
 Mar. NGC:006358
 New York State Department of Health - State/Local Government Agency [U.S.]. View all guidelines by the developer(s)</p> | <p>Compare</p> <p>Selected <input type="checkbox"/></p> <table border="1" style="width: 100%; border-collapse: collapse;"> <tr><td style="text-align: center;">12276</td></tr> <tr><td style="text-align: center;">0</td></tr> </table> | 12276 | 0 |
| 12276 | | | |
| 0 | | | |
| <p>10. Bipolar affective disorder. A national clinical guideline. 2005 May (reaffirmed 2008). NGC:004338
 Scottish Intercollegiate Guidelines Network - National Government Agency [Non-U.S.]. View all guidelines by the developer(s)</p> | <p>Compare</p> <p>Selected <input type="checkbox"/></p> <table border="1" style="width: 100%; border-collapse: collapse;"> <tr><td style="text-align: center;">7285</td></tr> <tr><td style="text-align: center;">0</td></tr> </table> | 7285 | 0 |
| 7285 | | | |
| 0 | | | |
| <p>11. Primary care diagnosis and management of adults with depression. 2004 Jan (revised 2012 Jan). NGC:009007
 Michigan Quality Improvement Consortium - Professional Association. View all guidelines by the developer(s)</p> | <p>Compare</p> <p>Selected <input type="checkbox"/></p> <table border="1" style="width: 100%; border-collapse: collapse;"> <tr><td style="text-align: center;">7285</td></tr> <tr><td style="text-align: center;">0</td></tr> </table> | 7285 | 0 |
| 7285 | | | |
| 0 | | | |

SIGN
1 fund

1)

BIPOLAR AFFECTIVE DISORDER

[Home](#)[Guidelines](#)
[Published Guidelines - by topic](#)[Published Guidelines - numbered list](#)[Rockets](#)[Audit tools](#)[SIGN Apps](#)[Suggest a Guideline Topic](#)[Current Work Programme](#)
[Consultation](#)[Events](#)[Implementation](#)[Patient Involvement](#)[Trainees](#)[Methodology](#)[Search](#)[About SIGN](#)

Scottish Intercollegiate Guidelines Network,
Healthcare Improvement Scotland, **Gyle Square, 1 South Gyle Crescent, Edinburgh EH12 9EB**
Tel. 0131 623 4720
Web contact duncan.service@nhs.net
Last modified 15/04/13
© SIGN 2001-2013

DOWNLOAD THE GUIDELINE

▶	Full guideline		(710K)
▶	Quick reference guide		(267K)
▶	Information about Acrobat files		

REMIT OF THE GUIDELINE

This guideline covers adults (aged 18 years or over) with bipolar affective disorders. A discussion of diagnostic issues in adolescence is included because of the often early onset of the condition.



HTA-databaseb

Søgt Bipolar disorder, afgrænset 2002-. Afgrænset til HTA in progress/ HTA Published



- **Search results [23 hits]**
- **Selected records [0 hits]**

Any field		AND	<input type="checkbox"/>	DARE	<input type="checkbox"/>	CRD assessed review (bibliographic)
Title	bipolar	OR	<input type="checkbox"/>	NHS	<input type="checkbox"/>	CRD assessed review (full abstract)
Author			<input type="checkbox"/>	EED	<input type="checkbox"/>	Cochrane review
Database entry date			<input type="checkbox"/>	HTA	<input type="checkbox"/>	Cochrane related review record
Publication year	2002 to 2013		<input checked="" type="checkbox"/>		<input type="checkbox"/>	CRD assessed economic evaluation (bibliographic)
			<input checked="" type="checkbox"/>		<input type="checkbox"/>	CRD assessed economic evaluation (full abstract)
					<input checked="" type="checkbox"/>	HTA in progress
					<input checked="" type="checkbox"/>	HTA published

Search now

Results for: ((bipolar):TI) and (Project record:ZDT OR Full publication record:ZDT) IN HTA FROM 2002 TO 2013

First	1	2	Last	Show all previews	Select all	Clear selections	Export
	Year	Database	Source	Title			
<input type="checkbox"/>	2011	HTA	HAYES, Inc.	Bipolar plasmakinetic electrovaporization for benign prostatic hyperplasia (BPH) [Preview]			
<input type="checkbox"/>	2011	HTA	HAYES, Inc.	Bipolar PlasmaKinetic electrovaporization for benign prostatic hyperplasia (BPH) [Preview]			
<input type="checkbox"/>	2011	HTA	Basque Office for Health Technology Assessment (OSTEBA)	[Case and control study: efficiency of the			

				psychoeducational treatment of the overload experienced by family members of bipolar patients] [Preview]	
<input type="checkbox"/>	2011	HTA	The Swedish Council on Health Technology Assessment (SBU)	[Long-term antipsychotic drug therapy for bipolar disorder] [Preview]	
<input type="checkbox"/>	2010	HTA	National Horizon Scanning Centre (NHSC)	Asenapine (Saphris) for manic episodes associated with bipolar I disorder [Preview]	
<input type="checkbox"/>	2010	HTA	Catalan Agency for Health Information, Assessment and Quality (CAHIAQ -formerly CAHTA)	[Guideline for the management of bipolar disorders in Catalonia] [Preview]	
<input type="checkbox"/>	2010	HTA	National Horizon Scanning Centre (NHSC)	Loxapine inhalation (Staccato loxapine) for agitation associated with schizophrenia or bipolar disorder [Preview]	
<input type="checkbox"/>	2010	HTA	HAYES, Inc.	Psychosocial interventions as adjunct treatments for pediatric bipolar disorder [Preview]	
<input type="checkbox"/>	2009	HTA	All Wales Therapeutics and Toxicology Centre (AWTTC), secretariat of the All Wales Medicines Strategy Group (AWMSG)	Aripiprazole (Abilify®) tablets for the treatment of moderate to severe manic episodes in Bipolar I Disorder and for the prevention of a new manic episode in patients who experienced	

				predominantly manic episodes and whose manic episodes responded to aripiprazole treatment [Preview]	
<input type="checkbox"/>	2009	HTA	National Horizon Scanning Centre (NHSC)	Risperidone long-acting injection (Risperdal Consta) for prevention of recurrent episodes in bipolar 1 disorder [Preview]	
<input type="checkbox"/>	2009	HTA	Unidad de Evaluacion de Tecnologias Sanitarias (UETS)	[Standards for the appropriate use of health technologies. Pharmacological treatment of bipolar disorder] [Preview]	
<input type="checkbox"/>	2008	HTA	Unidad de Evaluacion de Tecnologias Sanitarias (UETS)	[Appropriateness use standards for health technologies. Efficacy and safety of pharmacological treatment for bipolar disorder] [Preview]	
<input type="checkbox"/>	2008	HTA	National Horizon Scanning Centre (NHSC)	Aripiprazole (Abilify) for children and adolescents with bipolar disorder [Preview]	
<input type="checkbox"/>	2008	HTA	HAYES, Inc.	Medications for pediatric bipolar disorder: anticonvulsants [Preview]	
<input type="checkbox"/>	2008	HTA	HAYES, Inc.	Medications for pediatric bipolar disorder: atypical antipsychotics [Preview]	

<input type="checkbox"/>	2007	HTA	NIHR Health Technology Assessment programme	A systematic review and economic model of the clinical effectiveness and cost-effectiveness of interventions for preventing relapse in people with bipolar disorder [Preview]
--------------------------	------	-----	---	--

<input type="checkbox"/>	2006	HTA	National Institute for Health and Clinical Excellence (NICE)	Bipolar disorder: the management of bipolar disorder in adults, children and adolescents, in primary and secondary care [Preview]
--------------------------	------	-----	--	--

<input type="checkbox"/>	2005	HTA	NIHR Health Technology Assessment programme	Displaced intracapsular hip fractures in fit, older people: a randomised comparison of reduction and fixation, bipolar hemiarthroplasty and total hip arthroplasty [Preview]
--------------------------	------	-----	---	---

<input type="checkbox"/>	2004	HTA	NIHR Health Technology Assessment programme	A rapid and systematic review and economic evaluation of the clinical and cost-effectiveness of newer drugs for treatment of mania associated with bipolar affective disorder [Preview]
--------------------------	------	-----	---	--

<input type="checkbox"/>	2004	HTA	National Institute for Clinical Excellence (NICE)	Impedance-controlled bipolar radiofrequency ablation for menorrhagia [Preview]
--------------------------	------	-----	---	---

SBU, Sverige

Søgt Bipolär sjukdom, bipolar disorders

3 fund –1 rapport, 1 kommentar og 1 pågående

1)

Diagnostik och uppföljning av förstämningssyndrom

Slutsatser

- Det kliniska samtalet räcker inte för att diagnostisera depression och bipolära syndrom. Tillstånden upptäcks ofta inte. När bedömningen kompletteras med hjälpmedel i form av strukturerade eller semistrukturerade intervjuformulär och tillgång till journalhandlingar ökar den diagnostiska känsligheten (sensitivitet) och träffsäkerheten (specificitet).
- Två intervjuformulär, som täcker en hel rad psykiatriska diagnoser, har vetenskapligt stöd för en tillräckligt hög känslighet och träffsäkerhet för depression och mani hos vuxna. Det är den semistrukturerade Structured Clinical Interview for DSM-IV Axis I disorders (SCID-I) och den strukturerade Mini International Neuropsychiatric Interview (MINI). Den strukturerade intervjun Primary Care Evaluation of Mental Disorders (PRIME-MD) har för låg känslighet.
- Vid riktad screening av depression har följande formulär vetenskapligt stöd för tillräcklig känslighet: Child Behaviour Check List (CBCL) och Beck Depression Inventory-I (BDI-I) för barn och ungdomar, Beck Depression Inventory-II (BDI-II) för vuxna samt Geriatric Depression Scale (GDS-15 och GDS-30) för äldre utan påtaglig kognitiv svikt. Screening av nyförlösta mödrar med Edinburgh Postnatal Depression Scale (EPDS) identifierar cirka två tredjedelar av dem som har depression, vilket är i underkant. Formulären Patient Health Questionnaire-9 (PHQ-9) och Hospital Anxiety and Depression Scale (HADS) har för odefinierbar känslighet för att vara användbara.
Young Mania Rating Scale (YMRS) och General Behavior Inventory (GBI) för barn och ungdomar samt Hypomania Check List-32 (HCL-32) respektive Mood Disorder Questionnaire (MDQ med modifierade krav på funktionsnedsättning) för vuxna har vetenskapligt stöd för riktad screening av bipolärt syndrom.
- Det går inte att bedöma tillförlitligheten hos formulär som mäter svårighetsgrad av depression och mani. Det finns för få studier med tillräcklig kvalitet.
- Det går inte att bedöma tillförlitligheten hos formulär som är avsedda att följa förlopp av depression och mani. Det saknas studier.
- Det saknas studier som utvärderar kostnadseffektiviteten av att använda bedömningsformulär för förstämningssyndrom.
- Det finns många kunskapsluckor. Det viktigaste forskningsbehovet är att utvärdera hur strukturerad diagnostik som del i en systematisk handläggning påverkar tillfrisknande, relation mellan patient och behandlare samt kostnadseffektivitet. Utvärdering av intervjuformulär för barn och ungdomar samt formulär för bedömning av

depression hos äldre med kognitiv svikt är också angeläget. Det behövs ny forskningsmetodik för att utvärdera formulär som syftar till att följa sjukdomsförlopp.



- Användning av bedömningsformulär för förstämningssyndrom förutsätter att patienten har tillräckliga språkkunskaper samt, vid självbedömning, tillräcklig läs- och skrivförmåga. Det innebär att personer från andra kulturer, med läs- och skrivsvårigheter eller nedsatt kognitiv förmåga kan bli feltolkade eller ha svårt att fylla i formulären.

Citera till denna SBU-rapport:

SBU. Diagnostik och uppföljning av förstämningssyndrom. En systematisk litteraturöversikt. Stockholm: Statens beredning för medicinsk utvärdering (SBU); 2012. SBU-rapport nr 212. ISBN 978-91-85413-52-2.

Kort adress till sidan: <http://www.sbu.se/forstamningssyndrom>

- [Tipsa om denna sida](#)

- 
- 
- 

Ladda ner PDF



[Hela rapporten](#)



[Sammanfattning och slutsatser](#)

2)

Behandling med antipsykotiska eller stämningsstabiliserande läkemedel vid akut mani



[Skriv ut hela kommentaren \(pdf\) »](#)

[Startsida](#) / [Pågående](#) / Långtidsbehandling med antipsykotiska läkemedel vid bipolär sjukdom

3)

Långtidsbehandling med antipsykotiska läkemedel vid bipolär sjukdom

Projektledare: Mikael Nilsson

Projektassistent: Anna A Granath

Kontaktperson: Mikael Nilsson

E-post till kontaktperson: nilsson@sbu.se

Planerat publiceringsdatum: Hösten 2013

Projektgrupp:

Sakkunniga: Eva Lindström, docent i psykiatri, Akademiska sjukhuset och Leif Lindström, professor emeritus, Akademiska sjukhuset, Uppsala

Biträdande projektledare: Malin Höistad

Hälsoekonom: Emelie Heintz

Litteratursökare: Hanna Olofsson

Bakgrund

Användningen av andra generationens antipsykotiska läkemedel vid behandling av skov av mani, blandepisoder och svårare depressioner hos personer med bipolär sjukdom är relativt nytt. Dessa läkemedel används alltmer under en längre tid och i återfallsförebyggande syfte som monoterapi eller i kombination med litium eller annan stämningsstabiliserare.

Syfte

Utvärderingen syftar till att besvara hur långtidsbehandling (minst 6 månader) med antipsykotiska läkemedel påverkar patienter med bipolär sjukdom.

Socialstyrelsen, Sverige

Bipolär sjukdom, bipolar disorders

1 fund

1)

Nationella riktlinjer för depression och ångestsyndrom 2010 innehåller rekommendationer om effektivt omhändertagande, diagnostik, uppföljning och olika behandlingar som KBT, antidepressiva läkemedel och ECT.

Sök

- Sök i riktlinjerna


Uppföljning

-  Indikatorer för vård vid depression och ångestsyndrom
- Bevakning och uppdatering



Ladda ner

- Ladda ner/beställ riktlinjerna
- Patientversion av riktlinjerna

Om riktlinjerna

- Frågor och svar om riktlinjerna
- Centrala rekommendationer
- Sammanfattning – centrala rekommendationer
- Områden i riktlinjerna
- Så har riktlinjerna tagits fram
-  Hantering av synpunkter på den preliminära versionen av riktlinjerna

Effekter av psykosociala insatser för personer med schizofreni eller bipolär sjukdom - En sammanställning av systematiska översikter

-  Ladda ner sida
- [Share on facebook](#)[Share on twitter](#)[Share on email](#)[Share on print](#)
- RSS 
- Ladda ner
- Beställ
- Visa varukorgen

Sammanfattning

Psykiatrireformen som infördes 1995 innebar ett tydligare ansvar för kommunerna för att personer med psykiska funktionshinder skulle få sina sociala behov tillgodosedda och att vårdbehoven skulle tillgodoses av kommunerna i samverkan med landstingen. Tio år efter reformen finns behov att undersöka vilka vård- och omsorgsinsatser som är effektiva och hur dessa insatser ska kunna samordnas på ett för vårdtagaren tillfredsställande sätt.

[Läs hela sammanfattningen](#)

Publiceringsår: 2006

Artikelnnummer: 2006-110-37

ISBN: 91-85482-88-9

Format: Utskrift

Antal sidor: 153


Helsedirektoratet, Norge

Bipolar lidelse

1 fund

1)

Nasjonal faglig retningslinje for utgreiing og behandling av bipolare lidingar



- Tema: Psykisk helse og rus
- Kategori: Nasjonale faglige retningslinjerUtgiver: Helsedirektoratet

Ansvarlig avdeling: psykisk helsevern og rus

År: 2012Antall sider: 124

- IS-nummer: IS-1925

Målgruppe: Retningslinjen retter seg mot alle som yter helsetjenester til personer med bipolar lidelse

[LesHTML-versjon](#)[Last ned pdf](#)

Bestill

Bipolare lidelser har svært alvorlige konsekvenser for pasientenes livskvalitet og funksjonsevne. Lidelsene er vanskelige å diagnostisere, og de er både under- og overdiagnostisert.

Mange pasienter får ikke den behandlingen de trenger. Det faglige tilbudet til pasientene varierer, og det er uenighet om hvordan lidelsene skal utredes, diagnostiseres og behandles. Praksisvariasjonen er ikke faglig velbegrunnet og nyere behandlingsmetoder har blitt tatt i bruk på

svakt empirisk grunnlag. Det er nødvendig å bedre tilbudet, særlig med hensyn til psykososial oppfølging, psykoterapi og farmakoterapi.

Bipolare lidelser er forbundet med overdødelighet både på grunn av somatiske lidelser, ulykker og selvmord. Mennesker med bipolare lidelser vil være i behov av tiltak og oppfølging fra primærhelsetjenesten og det psykiske helsevernet og ofte veksle mellom tjenestenivåene. Behovet for koordinert samhandling er stort.

Retningslinjen gir kunnskapsbaserte råd om utredning og behandling for å sikre likeverdig og helhetlig tilbud til pasienter, uavhengig av hvor i landet de bor. De retter seg mot alle som yter helsetjenester til personer med bipolare lidelser.

Kunnskapscentret, Norge

0 fund

Cochrane Library:

Søgt på Bipolar disorder

1.	<ul style="list-style-type: none">• Topiramate for acute affective episodes in bipolar disorder <p>Kamini Vasudev , Karine Macritchie , John Geddes , Stuart Watson and Allan H Young</p> <p>January 2009</p> <p>1. Review</p>
2	Oxcarbazepine in the maintenance treatment of bipolar disorder
	<p>Akshya Vasudev , Karine Macritchie , Stuart Watson , John Geddes and Allan H Young</p> <p>January 2009</p> <p>2. Review</p>
3	Tiagabine in the maintenance treatment of bipolar disorder
	<p>Akshya Vasudev , Karine Macritchie , Sanjay NK Rao , John Geddes and Allan H Young</p> <p>June 2012</p> <p>3. Review</p> <ul style="list-style-type: none">•
4	

	<p>Lithium for maintenance treatment of mood disorders</p> <p>Sally SA Burgess , John Geddes , Keith KE Hawton , Matthew J Taylor , Ellen Townsend , K Jamison and Guy Goodwin</p> <p>January 2009</p> <p>1. Review</p> <ul style="list-style-type: none"> opt <input type="checkbox"/>
6.	<p>Family interventions for bipolar disorder</p> <p>Luis Justo , Bernardo Garcia de Oliveira Soares and Helena Calil</p> <p>January 2009</p> <p>1. Review</p>
7.	<p>Valproic acid, valproate and divalproex in the maintenance treatment of bipolar disorder</p> <p>Karine Macritchie , John Geddes , Jan Scott , D R Haslam and Guy Goodwin</p> <p>January 2009</p> <p>2. Review</p>
8	<p>Omega-3 fatty acids for bipolar disorder</p> <p>Paul Montgomery and Alexandra J Richardson</p> <p>January 2009</p> <p>3. Review</p>
9	<p>Tiagabine for acute affective episodes in bipolar disorder</p> <p>Akshya Vasudev , Karine Macritchie , Sanjay K Rao , John Geddes and Allan H Young</p> <p>December 2012</p> <p>4. Ns</p>

10	<p>Interventions for helping people recognise early signs of recurrence in bipolar disorder</p> <p>Richard Morriss , Mohammad Amir Faizal , Ashley P Jones , Paula R Williamson , Catherine A Bolton and James P McCarthy</p> <p>January 2009</p> <p>6. Review</p>
11	<p>Oxcarbazepine for acute affective episodes in bipolar disorder</p> <p>Akshya Vasudev , Karine Macritchie , Kamini Vasudev , Stuart Watson , John Geddes and Allan H Young</p> <p>December 2011</p> <p>7. Review</p>
12	<p>Antiglucocorticoid treatments for mood disorders</p> <p>Peter Gallagher , Navdeep Malik , James Newham , Allan H Young , I Nicol Ferrier and Paul Mackin</p> <p>January 2009</p> <p>8. Review</p>
13	<p>Olanzapine in long-term treatment for bipolar disorder</p> <p>Andrea Cipriani , Jennifer M Rendell and John Geddes</p> <p>January 2009</p> <p>9. Review</p>
14	<p>Valproate for acute mood episodes in bipolar disorder</p> <p>Karine Macritchie , John Geddes , Jan Scott , D R Haslam , Mauricio Silva de Lima and Guy Goodwin</p>

	<p>10.</p>
15	<p>Risperidone in long-term treatment for bipolar disorder</p> <p>Jennifer M Rendell and John Geddes</p> <p>January 2009</p> <p>11. Review</p> <ul style="list-style-type: none"> option15 <input type="checkbox"/>
16	<p>Lithium versus antidepressants in the long-term treatment of unipolar affective disorder</p> <p>Andrea Cipriani , Katharine A Smith , Sally SA Burgess , Stuart M Carney , Guy Goodwin and John Geddes</p> <p>July 2009</p> <p>1. Review</p>
17	<p>Haloperidol alone or in combination for acute mania</p> <p>Andrea Cipriani , Jennifer M Rendell and John Geddes</p> <p>July 2009</p> <p>2. Review</p>
18	<p>Pharmacotherapy for bipolar disorder in older people</p> <p>Akshya Vasudev , Alan J Thomas and Heinz Grunze</p> <p>April 2013</p> <p>3. Protocol</p>
19	<p>Antenatal psychosocial assessment for reducing perinatal mental health morbidity</p> <p>Marie-Paule Austin , Susan R Priest and Elizabeth A Sullivan</p> <p>October 2008</p>

	4. Review
20	<p>Olanzapine alone or in combination for acute mania</p> <p>Jennifer M Rendell , Harm J Gijsman , Paul K Keck , Guy Goodwin and John Geddes</p> <p>January 2009</p>
21	<p>Risperidone alone or in combination for acute mania</p> <p>Jennifer M Rendell , Harm J Gijsman , Mark S Bauer , Guy Goodwin and John Geddes</p> <p>January 2006</p> <p>5. Review</p>
21	<p>Droperidol for acute psychosis</p> <p>John Rathbone , Simone L Mandriota-Carpenter and Sharon J Cure</p> <p>January 2004</p> <p>6. Review</p>
22	<p>Valproate for schizophrenia</p> <p>Christian Schwarz , Anja Volz , Chunbo Li and Stefan Leucht</p> <p>July 2008</p> <p>7. Review</p>
23	<p>Adjunctive antioxidants for bipolar disorder</p> <p>Pedro V S Magalhães , Olivia Dean , Ana Cristina Andreazza , Michael Berk and Flávio Kapczinski</p> <p>October 2012</p>

24	<p style="color: blue; text-decoration: underline;">Interventions to encourage uptake of cancer screening for people with severe mental illness</p> <p>Elizabeth Barley , Rohan Borschmann , Paul Walters and Andre Tylee</p> <p>February 2012</p> <p style="background-color: #0070C0; color: white; padding: 2px;">8. Protocol</p>
----	---

PubMed

Search	Add to builder	Query	Items found	Time
#27	Add	Search "Bipolar Disorder"[Mesh] OR "bipolar disorder" Filters: Practice Guideline; Publication date from 2002/01/01	34	08:53:37
#16	Add	Search "Bipolar Disorder"[Mesh] Filters: Practice Guideline; Publication date from 2002/01/01	32	08:53:03
#26	Add	Search ((#4 or #23) and (#2 or #22)) Filters: Publication date from 2002/01/01	227	08:49:20
#25	Add	Search ((#4 or #23) and (#2 or #22))	271	08:42:42
#23	Add	Search "practice guideline"	19554	08:42:42
#22	Add	Search "bipolar disorder"	33545	08:42:42
#15	Add	Search "Bipolar Disorder"[Mesh] Filters: Practice Guideline	42	06:39:21
#2	Add	Search "Bipolar Disorder"[Mesh]	29553	06:39:04
#4	Add	Search "Practice Guidelines as Topic"[Mesh]	72643	06:35:21

EMBASE

59

#7

#5 AND #6

29,398

#6

'practice guideline'/exp/mj AND [2002-2013]/py

16,089

#5

'bipolar disorder'/exp/mj AND [2002-2013]/py

PsycInfo

Set	Search	Databases	Results	Actions
S4	SU.EXACT.EXPLODE("Bipolar Disorder") AND "practice guideline"	PsycINFO	18°	Actions
S3	"practice guideline"	PsycINFO	392°	Actions
S2	SU.EXACT.EXPLODE("Bipolar Disorder")	PsycINFO	18113*	Actions
S1	bipolar disorder	PsycINFO	23304*	Actions

Cinahl

Search ID#	Search Terms	Search Options	Actions
<input type="checkbox"/> S5	S2 AND S4	Search modes - Boolean/Phrase	View Results (3) View Details Edit
<input type="checkbox"/> S4	(MH "Practice Guidelines")	Search modes - Boolean/Phrase	View Results (29,961) View Details Edit
<input type="checkbox"/> S3	(MH "Practice Guidelines")	Search modes - Boolean/Phrase	View Results (0) View Details Edit
<input type="checkbox"/> S2	(MH "Bipolar Disorder+")	Limiters - Published Date from: 20020101-; Publication Type: Practice Guidelines Search modes - Boolean/Phrase	View Results (11) View Details Edit
<input type="checkbox"/> S1	(MH "Bipolar Disorder+")	Search modes - Boolean/Phrase	View Results (3,808) View Details Edit