

PICO 6 for ALKO DOB FINAL

Review information

Authors

[Empty name]¹

¹[Empty affiliation]

Citation example: [Empty name]. PICO 6 for ALKO DOB FINAL. Cochrane Database of Systematic Reviews [Year], Issue [Issue].

Characteristics of studies

Characteristics of included studies

Barrowclough 2001

Methods	
Participants	
Interventions	
Outcomes	
Notes	

Risk of bias table

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	alone. Individual patients were allocated to each condition by a third party with no affiliation to the study who used a computer-generated randomization list stratified for sex and three types of substance use (alcohol alone, drugs alone, or drugs and alcohol) to ensure equal male-female and substance use representation in each arm of the trial.

Allocation concealment (selection bias)	Unclear risk	Not described
Blinding of participants and personnel (performance bias)	Unclear risk	Single-blinded
Blinding of outcome assessment (detection bias)	Low risk	The assessors were blind to treatment allocation; attempts to maintain their blindness included use of separate rooms and administrative procedures for project staff, multiple coding of treatment allocations, and requesting subjects not to disclose information about the treatment.
Incomplete outcome data (attrition bias)	Unclear risk	Not described
Selective reporting (reporting bias)	Low risk	
Other bias	Low risk	

Footnotes

Characteristics of excluded studies

Footnotes

Characteristics of studies awaiting classification

Footnotes

Characteristics of ongoing studies

Footnotes

References to studies

Included studies

Barrowclough 2001

[Empty]

Excluded studies


Data and analyses

1 Intervention (family intervention) vs Control (no family intervention)

Outcome or Subgroup	Studies	Participants	Statistical Method	Effect Estimate
1.1 Funktionsniveau, GAF (level of functioning, GAF)	1	32	Mean Difference (IV, Fixed, 95% CI)	10.28 [0.22, 20.34]
1.1.1 Længste follow-up, min. 3-6 mdr. (Longest follow-up, min. 3-6 months)	1	32	Mean Difference (IV, Fixed, 95% CI)	10.28 [0.22, 20.34]
1.2 Frafald	1	36	Risk Ratio (M-H, Fixed, 95% CI)	0.33 [0.04, 2.91]
1.4 Psykiske symptomer (PANSS total score)	1	32	Mean Difference (IV, Fixed, 95% CI)	-6.52 [-17.85, 4.81]
1.4.1 Længste follow-up, min. 3-6 mdr. (Longest follow-up, min. 3-6 months)	1	32	Mean Difference (IV, Fixed, 95% CI)	-6.52 [-17.85, 4.81]
1.5 Psykiske symptomer (Positive symptomter)	1	32	Mean Difference (IV, Fixed, 95% CI)	-2.72 [-6.27, 0.83]
1.5.1 Længste follow-up, min. 3-6 mdr. (Longest follow-up, min. 3-6 months)	1	32	Mean Difference (IV, Fixed, 95% CI)	-2.72 [-6.27, 0.83]
1.6 Psykiske symptomer (Negative symptomter)	1	32	Mean Difference (IV, Fixed, 95% CI)	-2.02 [-5.88, 1.84]
1.6.1 Længste follow-up, min. 3-6 mdr. (Longest follow-up, min. 3-6 months)	1	32	Mean Difference (IV, Fixed, 95% CI)	-2.02 [-5.88, 1.84]

Figures

Figure 1 (Analysis 1.1)




Forest plot of comparison: 1 Intervention (family intervention) vs Control (no family intervention), outcome: 1.1 Funktionsniveau, GAF (level of functioning, GAF).

Figure 2 (Analysis 1.2)

Barrowclough 2001	Random sequence generation (selection bias)	+
	Allocation concealment (selection bias)	?
	Blinding of participants and personnel (performance bias)	?
	Blinding of outcome assessment (detection bias)	+
	Incomplete outcome data (attrition bias)	?
	Selective reporting (reporting bias)	+
	Other bias	+


Risk of bias summary: review authors' judgements about each risk of bias item for each included study.

Figure 4 (Analysis 1.4)




Forest plot of comparison: 1 Intervention (family intervention) vs Control (no family intervention), outcome: 1.4 Psykiske symptomer (PANSS total score).

Figure 5 (Analysis 1.5)



Forest plot of comparison: 1 Intervention (family intervention) vs Control (no family intervention), outcome: 1.5 Psykiske symptomer (Positive symptom).

Figure 6 (Analysis 1.6)



Forest plot of comparison: 1 Intervention (family intervention) vs Control (no family intervention), outcome: 1.6 Psykiske symptomer (Negative symptom).